

Semileptonic B Decays from an NRQCD/D234 Action

J.Shigemitsu^a, S.Collins^b, C.T.H.Davies^b, J.Hein^c, R.R.Horgan^d, G.P.Lepage^e

^aPhysics Department, The Ohio State University, Columbus, OH 43210, USA.

^bDepartment of Physics & Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK.

^cDepartment of Physics & Astronomy, University of Edinburgh, Edinburgh, EH9 3JZ, UK.

^dD.A.M.T.P., CMS, Wilberforce Rd., Cambridge, England CB3 0WA, UK.

^eNewman Laboratory, Cornell University, Ithaca, NY 14853, USA.

Semileptonic B decays are studied on quenched anisotropic lattices using Symanzik improved glue, NRQCD heavy quark and D234 light quark actions. We employ constrained fits to extract ground state contributions to two- and three-point correlators. Results for the $B \rightarrow \pi, l\bar{\nu}$ decay form factors are compared with previous lattice results. We find that our systematic errors (excluding quenching errors) are dominated by chiral extrapolation uncertainties.

1. Introduction

Several quenched lattice studies of $B \rightarrow \pi, l\bar{\nu}$ semileptonic decays have been completed in recent years [1]. They represent the first steps towards providing experimentalists and phenomenologists with the hadronic matrix elements involving the b quark that are necessary to carry out precision consistency tests of the Standard Model. In this article we present further studies of form factor calculations in the quenched approximation. We work with glue and quark actions that are more highly improved than in previous lattice calculations, we employ anisotropic lattices to enhance signal to noise and use constrained fits [2] to extract groundstate contributions to correlators in a controlled way. Simulation details, including the precise form of the action employed, can be found in [3]. We work at 5 values of the light quark mass spanning $0.7 m_{\text{strange}} \leq m_q \leq 1.3 m_{\text{strange}}$ and at one value of the heavy quark mass tuned to the b quark mass using the B_s meson. Results are then extrapolated to the chiral limit.

2. Fitting Three-point Correlators

The starting point for a form factor calculation on the lattice is the three-point correlator,

$$C_\mu^{(3)}(\vec{p}_B, \vec{p}_\pi, t_B, t) = \sum_{\vec{x}} \sum_{\vec{y}} e^{-i\vec{p}_B \cdot \vec{x}} e^{i(\vec{p}_B - \vec{p}_\pi) \cdot \vec{y}} \langle 0 | \Phi_B(t_B, \vec{x}) V_\mu^L(t, \vec{y}) \Phi_\pi^\dagger(0) | 0 \rangle. \quad (1)$$

$\Phi_\pi^\dagger$ and $\Phi_B^\dagger$ are interpolating operators used to create the pion or B meson respectively. V_μ^L is the dimensionless Euclidean space lattice heavy-light vector current. The continuum Minkowski space V_μ is related to V_μ^L via

$$V_\mu = a_s^{-3} Z_{V_\mu} \xi_\mu V_\mu^L. \quad (2)$$

ξ_μ is the conversion factor between Minkowski and Euclidean space γ -matrices and Z_{V_μ} is the heavy-light current matching coefficient. We work with B mesons decaying at rest ($\vec{p}_B = 0$). t_B is held fixed and we vary t , the location of the current insertions. The quantity $\xi_\mu C_\mu^{(3)}$ is fit, using constrained (Bayesian) fits, to the form,

$$\xi_\mu C_\mu^{(3)}(p_\pi, t) = \sum_j^{N_B} \sum_l^{N_\pi} A_{jl} e^{-E_\pi^{(l)} t} e^{-E_B^{(j)} (t_B - t)} \quad (3)$$

Constrained fits allow for multi-exponential fits to a single correlator without loosing stability. One augments the conventional χ^2 with a term, χ_{prior}^2 , which prevents fit parameters that are not constrained by the data from taking on “unreasonable” unphysical values and thus destabilizing the fit.

$$\chi^2 \longrightarrow \chi_{aug}^2 \equiv \chi^2 + \chi_{prior}^2, \quad (4)$$

with

$$\chi_{prior}^2 \equiv \sum_j \frac{(\alpha_j - \tilde{\alpha}_j)^2}{\tilde{\sigma}_j^2}. \quad (5)$$

One can check that those parameters, such as groundstate energies and amplitudes, that are well determined by the data, are only minimally affected by χ_{prior}^2 . In Figs. 1 and 2 we show fit results for the groundstate amplitude A_{11} for $\langle V_0 \rangle$ as a function of the number of exponentials N_π . N_B is held fixed at 1 or 2 and fits are carried out to all data points between $t_{min} = 1$ and $t_{max} = 17 \sim 22$. The numbers below data points give the $\chi_{aug}^2/d.o.f.$ of the fits. One sees that fit results, including errors, stabilize once $N_\pi > 2$. The fancy stars give bootstrap results at fixed $N_\pi = 3$ which we used in our final analysis (see references [2,3] for a description of constrained bootstrap fits).

3. Results for Form Factors and Comparisons with Previous Work

The groundstate amplitudes $A_{11}(V_\mu^L)$ extracted from three-point correlators are straightforwardly related to the continuum matrix element of interest.

$$\langle \pi(p_\pi) | V^\mu | B \rangle = \frac{A_{11}(V_\mu^L)}{\sqrt{\xi_\pi^{(1)} \xi_B^{(1)}}} 2 \sqrt{E_\pi M_B} Z_{V_\mu} \quad (6)$$

$\xi_\pi^{(1)}$ and $\xi_B^{(1)}$ are fixed from $\pi - \pi$ and $B - B$ two-point correlators. Fig. 3 shows results for the form factors $f_+(q^2)$ and $f_0(q^2)$ with the light quark mass around the strange mass. Errors reflect only statistical errors. Fig. 4 shows results after chiral extrapolation to the physical final state pion. One sees that errors have increased considerably.




Figure 1. Fit results for the groundstate amplitude A_{11} from the $\langle V_0 \rangle$ threepoint correlator. The numbers below the data points give $\chi_{aug}^2/d.o.f.$. The pion momentum is zero.




Figure 2. Same as Fig. 1 for pion momentum (111) in units of $2\pi/(a_s L)$.




Figure 3. The form factors f_+ and f_0 for the light quark mass fixed at the strange quark mass. Only statistical errors are shown.




Figure 4. The form factors f_+ and f_0 after chiral extrapolation to the physical pion. Statistical and chiral extrapolation errors shown.




Figure 5. Comparison of form factors from different lattice groups. Some errors are omitted.

In Fig. 5 we compare our results with those of other collaborations. Despite large differences in sources of systematic errors, one sees general agreement between all groups, especially for the form factor f_+ . All calculations suffer from large errors. In most cases, in particular in the present study, systematic errors are dominated by the chiral extrapolation uncertainties.

Acknowledgements : This work was supported by DOE Grant DE-FG02-91ER40690 and by PPARC and NSF.

REFERENCES

1. A. El-Khadra *et al.* [Fermilab Collaboration], Phys. Rev. **D64**:014502 (2001); S. Aoki *et al.* [JLQCD Collaboration], Phys. Rev. **D64**:114505 (2001); A. Abada *et al.* [APE Collaboration], Nucl. Phys. **B619**, 565 (2001); K.C. Bowler *et al.* [UKQCD Collaboration], Phys. Lett. **B486** (2000) 111.
2. G.P. Lepage *et al.*, Nucl. Phys. **B**(Proc. Suppl.)**106**, 12 (2002).
3. J. Shigemitsu *et al.*; hep-lat/0207011.